



Agromet Advisory Bulletin for the District, Kasaragod

(Valid from 20.12.2025 to 24.12.2025)

(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode & India Meteorological Department)



Bulletin Number: Pilicode/ Ksd – 101/2025

Date: 19/12/2025

A. Weather Summary of preceding four days

Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
0.0	31.0 – 34.0	15.2 – 21.0	57 – 96	00 – 01

B. Weather forecast for next five days

Parameters	20-12-2025	21-12-2025	22-12-2025	23-12-2025	24-12-2025
Average Rainfall, mm	0	0	0	0	0
Max. Temp, °C	34	34	34	34	34
Min. Temp, °C	21	21	21	21	21
Max. Relative Humidity, %	70	70	70	70	70
Min. Relative Humidity, %	65	65	65	65	65
Wind speed, km/h	3	3	3	3	3
Wind direction, degrees	270	230	290	270	270
Total cloud cover, octa	3	3	3	3	3

C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories
General Condition	No Rainfall**		
	The sky will be partially cloudy. Atmospheric humidity will be normal. Low night temperature and high day temperature will be experienced. Hence there will be distinct difference between day time temperature and night temperature.		
Weather warning	No rainfall from December 20 to 24.		
Impacts	Damages and losses may occur for crops having slender stems like banana and vegetables. High temperature during daytime may cause high rate of evaporation of water from the soil. Direct exposure to sunlight may cause sunburn and injuries animals.		
General Recommendations	Give popping to all soft slender stemmed crops like banana, vegetables, climbers etc Provide mulching to crops to reduce water loss from soil. For mulching, the trashes used should be free of any pest and diseases. Avoid the trashes of the same species as mulch. This will help to prevent the multiplication of crop specific pest and disease causing organisms. Powdering the top soil using a secondary tillage implement and spreading it uniformly over the field, will help to conserve water for a long period of time in the fields. Adopt drip irrigation method for maximum water use efficiency. Provide shade net for vegetable crops and ensure irrigation. Do not keep animals under open conditions for a long time. Provide sufficient drinking water to them.		

Coconut	All stages	Basal rot 	Drench the basin with Hexaconazole (25 ml in 25 L of water). After one week apply lime @ 2 kg per palm. Apply Potash @ 1 kg per palm three weeks after lime application to improve the health of palm, in addition to the regular dose. Avoid flood irrigation to prevent the spread of pathogen.
Banana	Various stages of growth	Sigatoka leaf spot 	Cut and burn all affected leaves. Spray Tilt or Contaf (@ 2ml per litre)
Amaranthus	All stages	Leaf blight 	Avoid overhead irrigation. Irrigate gently the basal regions of the plants. If disease is severe, apply cow dung slurry supernatant (2%) as foliar spray and Pseudomonas fluorescens (2%) or Trichoderma viridae (2%) as soil application.
Cucurbit Vegetables	Planting	Downy mildew 	As a prophylactic measure apply 'Mancozeb' (@ 2g/l of water). If disease appeared, spray Akomin® (@3ml/L) on both surfaces of the leaves, thrice at 15 days interval. Remove and burn out the infested leaves.
Cucurbitaceous Vegetables	Flowering and fruit setting	Melon fruit flies 	Keep pheromone traps. Spray Malathion:- Mix 2 ml of Malathion in one litre of water. Dissolve 10g of jaggery in the solution to attract the flies. Spray this solution on lower sides of the leaves
Chilli	Growth phase	Late spot 	Spray Mancozeb or Saaf (2g/litre of water) or Potassium phosphonate (2ml/litre of water)
Mango	Flowering stage	Mango hoppers 	Spray Azadirachtin (@5ml/litre Azadirachtin 3000pmm)

**** Warning colour codes of rainfall (for disaster management)**

Warning (Take actions)	Alert (Be prepared)	Watch (Be updated)	No warning (No actions)
-------------------------------	----------------------------	---------------------------	--------------------------------