



Agromet Advisory Bulletin for the District, Malappuram

(Valid from 15.10.2025 to 19.10.2025)

(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode & India Meteorological Department)



Bulletin Number: Pilicode/Mpm-82/2025

Date: 14/10/2025

A. Weather Summary of preceding five days

Rainfall, mm	Max. temp., °C	Min. temp., °C	R. H., %	Wind speed, Km/h
73.8	31.0 – 32.7	23.6 – 25.3	71 – 93	04– 10


B. Weather forecast for next five days

Parameters	15-10-2025	16-10-2025	17-10-2025	18-10-2025	19-10-2025
Average Rainfall, mm	4	4	2	1	3
Max. Temp, °C	33	33	33	33	33
Min. Temp, °C	25	25	25	25	25
Max. Relative Humidity, %	93	93	93	93	93
Min. Relative Humidity, %	71	71	71	71	71
Wind speed, km/h	2	2	2	2	2
Wind direction, degrees	270	270	270	230	230
Total cloud cover, octa	8	8	8	8	8

C. Agrometeorological Advisories

Crop	Stages	Problems	Agro-meteorological advisories
General Condition	Light to Moderate Rainfall**		
	The sky will be partially or fully cloudy. High relative humidity will be experienced. The atmospheric temperature will be high.		
Weather warning	There will be light to moderate rainfalls from October 15 to 17. There will be heavy rainfalls on October 18 in isolated places.		
Impacts	Damaging and losses or deteriorating quality of the open drying agricultural produces like copra, rubber, cashew nuts, seeds, food materials for processing etc.. Damages and losses may occur for crops having slender stems like banana and vegetables. Breaking branches and or trees may occur. High temperature during daytime may cause high rate of evaporation of water from the soil.		
General Recommendations	Do not attempt to dry the products like rice, rubber, copra, pulses, vegetable seeds, etc directly under sun. Give popping to all soft slender stemmed crops like banana, vegetables, climbers etc. Clean the drainage channels in crop lowlands to enable proper drainage of excess water in case of exigencies if any. Do not shelter the cattle in houses with weak structures. Keep animal feeds, seeds, fertilizers etc. in termite free and moist free rooms on elevated platforms coated with wooden panels. Provide mulching to crops to reduce waterloss from soil.		

Rice	Transplanting stage for second crop	Dip the roots of the seedlings in slurry of pseudomonas (20g pseudomonas/litre of water) for 30 minutes just before transplanting. This will help the seedlings to grow vigorously. Apply fertilizers at the occasion of no rainfalls, after lowering the water and blocking all drainage channels. While transplanting, apply 400g urea, 900g mussooriephos and 150g muriate of potash for the medium duration varieties and 400g urea, 700g mussooriephos and 120g muriate of potash for the short duration varieties to each cent (40 square meter) of land area.	
Cool season vegetables	Land preparation	Incorporate lime or dolomite to the soil depending on its acidity, at least 14 days before the application of chemical fertilizers. This will not only reduce the acidity but also control the soil borne diseases.	
Coconut	All stages	Stem bleeding 	Chisel out the affected parts and apply Bordeaux paste. Drench the palm basin (2m radius) with 40 litres of 1% Bordeaux mixture or soil drenching of Hexaconazole (Contaf®) 25ml/25 litre of water once in 4 months. After one month, apply 5 kg Trichoderma enriched neem cake and 500g of Ayar® (micro nutrient mixture)
Areca nut	All stages	Mahali disease 	Apply Bordeaux mixture (1%) to the bunch and leaves.
Banana	Various stages of growth	Pitting disease 	Spray Mancozeb (@2g/litre), only on immature bunches. . Cover the bunches with poly ethylene sheets having pin holes on it. This will help to protect the bunches from birds attack also.
Ginger	All stages	Rhizoctonia leaf blight 	Spray carbendazim (@2g/litre of water)

Poultry	All stages of growth	Newcastle Disease 	Newcastle disease or Ranikhet disease is a highly contagious disease of birds caused by a para-myxo virus. This infection results gasping and coughing, drooping wings, dragging legs, twisting of the head and neck, circling, in appetite, complete paralysis. Prevention: Ensure complete hygiene condition for birds and chicken coops. Provide vaccinations with Live B1 and La Sota strains administered in drinking water intranasally or intraocularly.
---------	----------------------	--	---

**** Warning colour codes of rainfall (for disaster management)**

Warning (Take actions)	Alert (Be prepared)	Watch (Be updated)	No warning (No actions)
-------------------------------	----------------------------	---------------------------	--------------------------------

Sd/-
Nodal Officer,
GKMS Project, RARS Pilicode